

Cannabinoid profile of FINOLA inflorescences. The value for each listed cannabinoid is calculated from the average of five independent replicates.

Cannabinoid		Mean (µg/g)	Mean %
<i>Neutral forms</i>			
CBD	Cannabidiol	2,614	0.26
Δ9-THC	Tetrahydrocannabinol	299	0.03
CBN	Cannabinol	12	0.01
CBC	Cannabichromene	< 1 µg/g	< 0.00
CBG	Cannabigerol	190	0.02
CBDV	Cannabidivarin	4,888	0.49
Δ9-THCV	Tetrahydrocannabivarin	< 1 µg/g	< 0.00
<i>Acid forms</i>			
CBDA	Cannabidiolic acid	23,479	2.35
Δ9-THCA	Tetrahydrocannabinolic acid	384	0.04
CBNA	Cannabinolic acid	214	0.02
CBCA	Cannabichromenic acid	457	0.46
CBGA	Cannabigerolic acid	698	0.07
CBDVA	Cannabidivarinic acid	2,862	0.29
Δ9-THCVA	Tetrahydrocannabivarinic acid	< 1 µg/g	< 0.00

Source: Table 6, in “Phytochemical and Ecological Analysis of Two Varieties of Hemp (*Cannabis sativa* L.) Grown in a Mountain Environment of Italian Alps”, by Radmila Pavlovic, Sara Panseri, Luca Giupponi, Valeria Leoni, Cinzia Citti, Chiara Cattaneo, Maria Cavaletto and Annamaria Giorgi, published in *Frontiers in Plant Science*, October 2019, Volume 10, Article 1265, doi: 10.3389/fpls.2019.01265.